

# 秦岭六种提灯藓的染色体研究\*

肖娅萍 刘全宏 田先华

(陕西师范大学生物系, 西安 710062)

## CHROMOSOME NUMBERS OF SIX SPECIES OF *MNIUM* FROM QINLING RANGE IN CHINA

Xiao Ya-ping Liu Quan-hong Tian Xian-hua

(Department of Biology, Shaanxi Normal University, Xi'an 710062)

**Abstract** Reported in this paper are chromosome numbers of six species of *Mnium* from Qinling Range. The chromosome number of *M. pseudolycopodioides*,  $n=6$ , is reported here for the first time. The counts for *M. marginatum* and *M. lycopodioides*,  $n=12$  and  $n=6$  respectively, agree with the previous reports, and those for *M. laevinerve*, *M. heterophyllum* and *M. thomsonii*, here all found to be  $n=6$ , differs from some or all previously published chromosome numbers for these three species.

**Key words** *Mnium*; Mitosis; Meiosis; Chromosome numbers

**摘要** 本文以藓类植物的幼嫩孢蒴及茎尖为材料,对产于中国秦岭的六种提灯藓的染色体进行了观察计数。首次报道了 *M. pseudolycopodioides* 的染色体数目为  $n=6$ ; *M. marginatum* 和 *M. lycopodioides* 的染色体数分别为  $n=12$  及  $n=6$ , 与前人报道的结果一致; *M. Laevinerve*、*M. heterophyllum* 以及 *M. thomsonii* 的染色体数目均为  $n=6$ , 与前人的报道不完全相同。本文旨在为本属的研究提供一定的细胞学资料。

**关键词** 提灯藓属;有丝分裂;减数分裂;染色体数目

*Mnium* is a large genus in the family Mniaceae, consisting of about 100 species in the world. This genus has a global distribution but is most richly represented in the warm temperate regions of the Northern Hemisphere. In China, 49 species have been recorded, 29 of which are native to Qinling Range (Northwest Institute of Botany, 1978; Li *et al.*, 1979).

*Mnium* is the first moss genus to which cytological data were applied to solve its systematic problems (Lowry, 1948). Since then, numerous cytological and cytotaxonomical studies have been made on this genus (Bowers, 1980, 1968; Bryan, 1973; Ono, 1970, 1967a, 1967b, 1967c; Tatuno *et al.*, 1966; Holmen, 1958). Lowry (1948) and Bowers (1980, 1968) pointed out that cytological data can provide quite important information for the clarification of the intergeneric relationships of the family Mniaceae and the interspecific relationships within the genus *Mnium*.

In this paper, chromosome numbers of six species in *Mnium* from Qinling Range are reported, with a view to offering some cytological evidence for a better understanding of the interspecific relationships of *Mnium* in this area.

## 1 Materials and Methods

All the plants studied here were collected from Qinling Range (Table 1). The voucher specimens are deposited in the Herbarium of Department of Biology, Shaanxi Normal University (SANU).

Table 1 The origin of the experimental materials

Species	Habitat	Voucher
<i>M. pseudolycopodioides</i> C. Muell.	Sunny slope, 1300 m	Y. P. Xiao, W. G. Fu, 924002
<i>M. marginatum</i> (With.) P. Beauv.	Shadowy slope, 1400 m	Y. P. Xiao, Y. R. Liu, 923008
<i>M. lycopodioides</i> Schwaegr.	Shadowy slope, 1400 m	Y. P. Xiao, 923009
<i>M. laevinerve</i> Card.	Sunny slope, 1900 m	Y. P. Xiao, 924032
<i>M. heterophyllum</i> (Hook.) Schwaegr.	Sunny slope, 1300 m	Y. P. Xiao, Y. R. Liu, 924001
<i>M. thomsonii</i> Schimp.	Shadowy slope, 1500 m	Y. P. Xiao, Y. R. Liu, 923038

The technique for chromosome preparation follows Zhou *et al.* (1988). In the field, immature capsules and stem apices were fixed in Carnoy's fluid (absolute alcohol: glacial acetic acid = 3:1) for 4~12 hours, and then transferred to 70% alcohol and kept in it at low temperature. When making preparations of sporocyte meiosis, calyptra and operculum were removed in Carnoy's fluid with scalpel under stereomicroscope, and the sporocyte layer was then stripped carefully from the surface of columella, and stained in carbol fuchsin. When making preparations of stem apex mitosis, the stem apices were not treated and were directly placed on slides, on which a drop of carbol fuchsin fluid was added for staining. For permanent preparation, the slides and coverslips were separated rapidly after being frozen with liquid nitrogen and the slides were washed with absolute ethanol for 2~3 min. The preparations were finally remounted in gum arabic. The chromosomes were observed and photomicrographed under an Olympus BH2 microscope.

## 2 Results

### 2.1 *M. pseudolycopodioides* C. Muell.

Six bivalents were observed at metaphase I of sporocyte meiosis (Plate 1:1). Thus, the chromosome number of this species is  $n=6$ , which is reported here for the first time.

### 2.2 *M. marginatum* (With.) P. Beauv.

Twelve chromosomes were observed at mitotic metaphase of stem apices of the gametophytes (Plate 1:2). This species has been cytologically studied previously by Lowry (1948), Holmen (1958), Bowers (1980,1968), Smith *et al.* (1968), Bryan (1973), Kumar *et al.* (1983). All obtained the same result.

### 2.3 *M. lycopodioides* Schwaegr.

From our observations on the metaphase I and anaphase I of sporocyte meiosis (Plate 1; 3,4), the chromosome number of this species was counted as  $n=6$ . The result agrees with the previous reports by Ono *et al.* (1977) and Kumar *et al.* (1987). The plants studied by Bowers (1980) under the name *M. lycopodioides*, which have also the number of  $n=6$ , however, should be actually identified as *M. ambiguum* (Koponen, 1981).

### 2.4 *M. laevinerve* Card.

Our observations on the metaphase I and anaphase I of sporocyte meiosis (Plate 1; 5,6) showed that this species had the chromosome number of  $n=6$ . The result agrees with the previous report by Kumar (1973), but disagrees with the reports of  $n=7$  by Tatuno *et al.* (1966) and by Ono (1970) for this species.

### 2.5 *M. heterophyllum* (Hook.) Schwaegr.

Our observations on the metaphase I and anaphase I of sporocyte meiosis (Plate 1; 7,8) showed that this species had the chromosome number of  $n=6$ . Ono (1970; 1967c, under the name *M. sapporensis*) twice reported the chromosome number of this species as  $n=7$ , while Kumar *et al.* (1983) reported it as  $n=12$ , but Kumar *et al.* (1987) reported it as  $n=6$ .

### 2.6 *M. thomsonii* Schimp.

Our observation on the metaphase I and anaphase I of sporocyte meiosis (Plate 1; 9,10) showed that this species had the chromosome number of  $n=6$ . The result agrees with the previous report by Bowers (1968) but disagrees with the report of  $n=8$  by Ono (1970) for this species. Kuta *et al.* (1984) reported two chromosome numbers of  $n=6$  and 12 for this species from Poland.

## 3 Discussions

Lowry (1948) proposed the basic chromosome number of the genus *Mnium* as  $n=6$ , but Tatuno *et al.* (1966) established it as  $n=7$ . All the six species studied here have the basic number of  $n=6$ .

The genus *Mnium* can be conveniently divided into three groups based on the marginal teeth of the leaves, i. e. single-toothed group, double-toothed group and entire-leaved group. This grouping, as Bowers (1968) pointed out, is not so artificial as it would seem at first glance since it is supported by cytological evidence. Bowers (1968) found that all the entire-leaved species he studied have the basic chromosome number of  $n=6$ , and that although the cytological evidence separating the single-toothed species from the double-toothed species is not too convincing because of the same basic chromosome number of  $n=6$  in both groups, they can still be cytologically separated on the basis of the presence of subterminal or terminal spindle-attachment regions in the double-toothed species and the absence of these in the single-toothed species. All the six species studied here belong to the double-toothed group and have the basic chromosome number of  $n=6$ . Thus, our results, though without

reference to the chromosome morphology, at least to some extent support Bower's generalization.

*Mnium marginatum* is the only synoicous one of the six species studied, and it has the chromosome number of  $n=12$ . In the genus *Mnium*, a diploid gametophyte number is usually correlated with the synoicous condition. Our result reinforces this point. As regards to its origin, Bowers (1968) considered that *M. marginatum* might be an autodiploid derived from *M. ambiguum*.

Our count of  $n=6$  for *M. laevinerve* agrees with the previous report by Kumar (1973) but disagrees with the reports of  $n=7$  by Tatuno *et al.* (1966) and Ono (1970). In the latter two authors' papers, there are clearly seven chromosomes in this species which consist of six long and one very short, or "m" chromosomes. In fact, almost all the species of *Mnium* studied by these two authors have the chromosome number of  $n=7$ . Both of them attached great importance to the observation of the "m" chromosome, which was rarely reported by other authors.

More extensive and intensive cytological studies, especially those with proper reference to gross morphological characters and sexual reproduction organism, should be carried out for a further understanding of the interspecific relationships within the genus *Mnium* in the Qinling Range.

## References

- Bowers M C. 1968. A cytotaxonomic study of the genus *Mnium* in Colorado. *Rev Bryol Lichenol.* **36**: 167~202
- Bowers M C. 1980. A cytotaxonomic classification of the Mniaceae (Bryophyta). *Lindbergia.* **6**: 22~31
- Bryan V S. 1973. Chromosome studies on Mosses from Austria, Czechoslovakia and other part of central Europe. *österr Bot Zeit.* **121**: 187~226
- Holmen K. 1958. Cytotaxonomic studies in some Danish mosses. *Bot Tids.* **54**: 23~43
- Koponen T. 1981 Miscellaneous notes on Mniaceae (Bryophyta) VII. Verification of the vouchers of some chromosome number reports. *Ann Bot Fennici.* **18**: 73~82
- Kumar S S. 1973. Cytological studies on some west Himalayan mosses I. *The Bryologist.* **76**: 202~206
- kumar S S. Garg B K. 1983. Chromosome numbers in some west Himalayan mosses. *Misc Bryol Lich.* **9**: 171~172
- Kumar S S. Koponen T. Uniyal P L. 1987. In: Love A ed. Chromosome number reports XC VII. *Taxon.* **33**: 766~767
- Kuta E. Przywara L. Ochyra R. 1984. Chromosome studies on polish bryophytes I. *Beitr.* **3**: 28~45
- Li X J. Zang M. 1979. Studies on Chinese Mniaceae. *Acta Bot Yunn.* **1**: 32~80
- Lowry R J. 1948. A cytotaxonomic study of the genus *Mnium*. *Mem Torrey Bot Club.* **20**: 1~42
- Northwest Institute of Botany. Academia Sinica. 1978. Bryophyta. *Flora Tsinglingensis*. Vol 3. Beijing: Science Press
- Ono K. 1967a. Structural sex chromosomes of Mniaceae I. *Bot Mag (Tokyo).* **80**: 185~191
- Ono K. 1967b. Structural sex chromosomes of Mniaceae I. *Bot Mag (Tokyo).* **80**: 363~372
- Ono K. 1967c. Structural sex chromosomes of Mniaceae III. *Bot Mag (Tokyo).* **80**: 498~507
- Ono K. 1970. Karyological studies on Mniaceae and Polytrichaceae. with special reference to the structural sex chromosomes 2. *J Sci Hiroshima Univ. Ser B. Div 2 (Bot)* **13**: 107~166

- Ono K., Morita Y., Taguchi M. *et al.*, 1977. On the chromosome number in Mniaceae. *Kumamoto J Sci Biol.* **13**: 49~65
- Smith A J E., Newton M E., 1968. Chromosome studies on some British and Irish mosses **II**. *Trans Brit Bryol Soc.* **5**: 463~522
- Tatuno S., Ono K., 1966. Zytologische Untersuchungen über die Arten von Mniaceae aus Japan. *J Hattori Bot Lab.* **29**: 79~95
- Zhou Y L., Sha W., Wang C K *et al.*, 1988. Cytological observation of 4 Chinese moss species. *Acta Phytotax Sinica.* **26**: 378~381

### Explanation of Plate


**Plate 1** 1,3~10. Meiosis in sporocytes; 2. Mitosis in stem apices.

1. *M. pseudolycopodioides* C. Muell.,  $n=6$ , 1. MI; 2. *M. marginatum* (With) P. Beauv.,  $n=12$ ; 3~4. *M. lycopodioides* Schwaegr.,  $n=6$ , 3. MI, 4. AI; 5~6. *M. laevinerve* Card.,  $n=6$ , 5. MI, 6. AI; 7~8. *M. heterophyllum* (Hook.) Schwaegr.,  $n=6$ , 7. MI, 8. AI; 9~10. *M. thomsonii* Schimp.,  $n=6$ , 9. MI, 10. AI.

(上接 287 页)

- 6 1992 年国家科学技术进步奖一等奖(获奖项目的一部分)  
1991 年国家中医药管理局中医药科技进步奖二等奖(获奖项目一部分)  
乔传卓,朱友平,苏中武,李承祜:杏属一新种及其近缘类群的修订。31(2) (1992)
- 7 1994 年浙江省人民政府颁发的科学技术进步奖一等奖(获奖项目的一部分)  
张渝华:关于浙江泡果芥和棒毛芥的分类问题。33(2) (1995)  
张渝华:安徽泡果芥属(十字花科)一新种。33(1) (1995)
- 8 1994 年中国科学院自然科学奖二等奖(获奖项目的一部分)  
李承森:徐氏蕨属—中国云南早泥盆世陆地植物的一个新属。20(3)  
耿宝印:四川早泥盆世植物群的研究。30(3) (1992)
- 9 1994 年中国科学院自然科学奖三等奖(获奖项目的一部分)  
何关福,马忠武,印万芬,陈梦玲:银杉的锯齿稀族成分与系统位置的讨论。19(4)
- 10 1993 新疆维吾尔自治区科学技术进步奖二等奖(获奖项目的一部分)  
蓝永珍,周太炎:中国芸苔属的新资料。29(1) (1991)
- 11 1995 年中国科学院自然科学奖三等奖(获奖项目一部分)  
吴鹏程,李登科,高彩华:武夷山苔藓植物区系及其与邻近地区的关系。25(5)  
吴鹏程,罗健馨,汪楣芝:一个原始的苔类的目—藓苔目在中国的发现。21(1)  
吴鹏程:西藏苔藓植物两新种。20(3)  
吴鹏程,罗健馨:东喜马拉雅南翼苔藓植物区系的特性及其来源。20(4)  
陈邦杰,吴鹏程:中国羽苔属一新种。17(2)  
吴鹏程,林邦娟:海南苔类植物区系与地理分布的初步讨论。16(2)  
吴鹏程,罗健馨,孟繁松:第 4 纪藓类化石在中国的发现。14(1)  
胡晓云,吴鹏程:四川金佛山藓类植物区系的研究。29(4) (1991)
- 12 1995 年安徽省高等学校科学技术进步奖二等奖(获奖项目的一部分)  
1991—1993 年安徽省自然科学优秀论文三等奖  
郭新弧:国产唇形科新分类群。31(3) (1993)
- 13 1995 年安徽省高等学校科学技术进步奖二等奖(获奖项目的一部分)  
张定成,邵建章:安徽百合科二新种。29(5) (1991)  
张定成,邵建章,汪小全:安徽南部贝母属植物核型研究。30(1) (1992)

(下转 326 页)



See explanation at the end of text